

NEMASYS GROW YOUR OWN PACK

Pest	Equipment/ How to Apply	Area Covered	When to Apply	Frequency	Things to Note	Pest
Carrot Root Fly	Use a watering can with a coarse rose or a hose end feeder. Apply as a drench around roots of the plants or where pest is present. Apply to moist soil and water in.	Up to 60m ²	April to July. Apply after planting and repeat fortnightly throughout the growing season when pests are present.	Fortnightly application maintains control and catches subsequent generations.	Apply along the rows and around the plants. The pests are next to the stems and roots of plants.	Carrot Root Fly
Cabbage Root Fly						Cabbage Root Fly
Cutworm						Cutworm
Onion Fly						Onion Fly
Leatherjacket				Repeat applications are required to control the pest.	Apply to the whole soil area.	Leatherjacket
Sciarid					Greenhouse pest. Apply to the soil in pots, growing bags or open soil.	Sciarid
Ants					Ants do not tolerate the nematodes near their nests and move the nests away from these areas. Apply to the soil area or to individual ant nests.	Ants
Caterpillars	Use a pump sprayer. Spray the pests on the plants. Pests not directly sprayed will not be controlled. Wet areas before applying. Avoid direct sunlight.	Up to 60m ²	Apply when pest is present. Consider weekly applications until all pests have hatched out.	3 applications are recommended to control all pests hatching out.	Contact with pest is essential.	Caterpillars
Gooseberry Sawfly						Gooseberry Sawfly
Thrips						Thrips
Codling Moth	Use a pump sprayer. Apply to the trunk, main branches and soil beneath the canopy. Wet areas before applying.	Up to 16 Trees	Apply during September or October. Make 3 weekly applications to control all pests hatching out.	3 applications are essential to control all pests hatching out.	Provides control of the overwintering stages and therefore protection for the following year.	Codling Moth

NB: Timings will vary depending upon location in the country and whether treating pests outside or undercover